

ORIGINAL RESEARCH

Open Access

# OX40 ligand expression abrogates the immunosuppressive function of retinal pigment epithelium

Matthew A Cunningham<sup>1\*</sup>, Zhuqing Li<sup>2</sup>, Baoying Liu<sup>3</sup>, Steven Yeh<sup>4</sup> and Robert B Nussenblatt<sup>3</sup>

## Abstract

**Background:** This study aims to investigate the role of OX40 ligand (OX40L) in ocular inflammation via abrogation of retinal pigment epithelium (RPE)-mediated immunosuppression using an *in vitro* expression approach. OX40L cDNA was polymerase chain reaction-amplified and cloned into an eYFP fusion vector. Cultured retinal pigment epithelial cells (ARPE-19) were transfected with the vector. Total RNA from unstimulated or inflammatory cytokine-stimulated ARPE cells were isolated and analyzed for OX40L expression by reverse transcription-polymerase chain reaction. Peripheral blood mononuclear cells (PBMCs) were isolated from healthy human donors. Human ARPE cells ( $\pm$ OX40L  $\pm$  GITR ligand (GITRL) expression) and PBMCs were co-cultured for *in vitro* proliferation studies.

**Results:** Polymerase chain reaction confirmed the insertion of the OX40L gene into the fusion vector. Flow cytometry and fluorescence microscopy further confirmed surface expression of OX40L on ARPE cells after transfection. OX40L expression was induced in the RPE cells stimulated with pro-inflammatory cytokines. In the co-culture studies, there was a significant reversal (20% to 30%) of the RPE-induced suppression of activated PBMCs when the ARPE cells were transfected with OX40L. When both OX40L and GITRL were concomitantly transfected into ARPE cells, there was an additive reversal of RPE-mediated T cell suppression, when compared to the reversal caused by RPE cells expressing either OX40L alone or GITRL alone.

**Conclusions:** Using an *in vitro* approach, we found that OX40L causes an abrogation of the RPE-mediated immunosuppression. OX40L appears to be regulated in the ARPE-19 cell line and may play an important role in the pathogenesis of various ocular inflammatory conditions.

**Keywords:** OX40 ligand, Uveitis, Retinal pigment epithelium, Tumor necrosis factor ligand, *in vitro*

## Background

OX40 ligand (OX40L) is a member within the TNF ligand superfamily [1-4] and plays an important role in the induction of co-stimulatory signaling during T cell and antigen-presenting cell (APC) interactions [5-9]. Prior studies have demonstrated that OX40L is expressed predominantly on CD4<sup>+</sup> T cells [10,11]. In addition to its role in co-stimulatory signaling, OX40/OX40L interaction is closely involved in effector functions [12-15]. OX40L expression has been observed on APCs including dendritic cells [16,17], B cells [12], and microglial

cells [18], but has also been found on non-lymphoid cells (i.e., endothelial cells) [19].

Recently, its potential role in the pathogenesis of immune-mediated ophthalmic diseases including uveitis, experimental allergic conjunctivitis, and corneal graft rejection has been investigated [20-23]. A recent study by Zhang et al. described the activation of ovalbumin-specific T cells with OX40-activating antibody and found that OX40-stimulated lymphocytes elicited more ocular inflammation compared to donor cells without OX40 activation. Moreover, flow cytometry showed that Th17 cells expressed OX40 ligand, and interleukin-17 neutralizing antibody was successful in the attenuation of OX40-mediated uveitis [22]. In another study of experimental autoimmune uveitis, OX40 activation was found to promote the expansion uveitogenic memory T cells, as

\* Correspondence: macunning@gmail.com

<sup>1</sup>Vitreoretinal Service, Department of Ophthalmology and Visual Sciences, The University of Iowa Hospitals & Clinics, Iowa City, IA 52242, USA  
Full list of author information is available at the end of the article

evidenced by the upregulation of IL-7R in CD4<sup>+</sup>CD44<sup>+</sup> lymphocytes [23].

Other animal models of ocular inflammation have been described. In a murine model of experimental allergic conjunctivitis, *in vivo* stimulation of OX40 with an anti-OX40 antibody led to the exacerbation of ocular surface inflammatory disease, while antibody blockade of the OX40/OX40L interaction inhibited disease expression [20]. Hattori et al. characterized OX40/OX40L interaction in a murine model of corneal transplant rejection. They describe significantly reduced corneal allograft rejection in mice treated with a monoclonal antibody targeting murine OX40L when compared to control mice. In addition, OX40 ligand-deficient mice demonstrated a decreased risk of allograft rejection when transplanted with wild-type donor corneas. An *in vitro* study also revealed that treatment with anti-OX40L monoclonal antibody reduced the proliferative response and interferon-gamma production of lymph node cells after donor alloantigen stimulation. These studies were suggestive that OX40L blockade prolonged graft survival via inhibition of recipient T cell activation [21].

Thus, while OX40L and OX40 in activating inflammation has been suggested by several animal models of systemic and ophthalmic inflammation, the precise pathways by which OX40L leads to these ocular inflammatory sequelae warrants further investigation. The eye is an immune-privileged organ, and multiple mechanisms are thought to contribute to this privilege [24-26] including retinal pigment epithelium (RPE)-mediated suppression of T cell activation [27], induction of apoptosis of activated T cells via Fas-FasL interactions [28], and inhibition of T cell proliferation by a cell-contact-dependent mechanism [29]. Using an *in vitro* expression system previously described in a report for GITR ligand [30], another important member of the TNF superfamily, we sought to further investigate the role of OX40L and its role in ocular inflammation, specifically in its ability to abrogate RPE-mediated immunosuppression. We found that OX40L causes abrogation of RPE-mediated immunosuppression and appears to be regulated within RPE cells.


## Results

### Confirmation of OX40L insertion into eYFP vector

To confirm the correct insertion of the OX40L gene into the eYFP vector, polymerase chain reaction (PCR) was performed on the purified products. OX40L was detected using PCR (Figure 1), along with restriction enzyme (BamH1 and HindIII) overnight digestion. The gene was identified in both the eYFP-C1 and eYFP-N1 vectors.


### Expression of OX40L on ARPE-19 cells

Using flow cytometry and a biotinylated anti-OX40L antibody, we found that the expression of OX40L was



undetectable (data not shown). After transfection, OX40L was expressed on RPE cells, and the expression was monitored using both flow cytometry and fluorescence microscopy.

In a time and dosage trial, after verified transfection of the eYFP-OX40L construct into RPE cells, surface expression of OX40L was demonstrated as early as 4 h after transfection (see Figure 2). The expression of our protein was primarily cytosolic prior to the 4-h mark. This was



demonstrated by the flow cytometric detection of the eYFP fluorescent protein as early as 30 min after transfection. During the dosage trial, transfecting a dose of 0.5  $\mu$ g per  $0.6 \times 10^6$  RPE cells led to an expression >90% (data not shown). Larger transfection dosages led to a plateau effect, and dosages >5  $\mu$ g, in the eYFP-OX40L-C1 construct, had a toxic effect to the RPE cells (data not shown).

#### Cytokine-mediated regulation of OX40L RNA in cultured RPE

Using reverse transcription-polymerase chain reaction (RT-PCR), the OX40L RNA content in resting, unstimulated RPE cells and in RPE cells stimulated for 24 h with high levels of the pro-inflammatory cytokines IL-1 $\alpha$ , TNF- $\alpha$ , and IFN- $\gamma$  were compared. Constitutive expression of OX40L RNA was not observed in unstimulated RPE cells; however, in the stimulated RPE cells, OX40L was expressed (Figure 3).


#### Transient expression of OX40L in RPE cell reverse RPE-induced suppression of PBMCs

A co-culture of RPE cells (with or without OX40L expression) and PBMCs (with or without anti-CD3/anti-

CD28 stimulation) was performed. RPE cells without OX40L expression, when co-cultured with either stimulated or unstimulated PBMCs, demonstrated the greatest suppressive activity (Figure 4). When RPE cells were transfected with either the eYFP-C1-OX40L or the eYFP-N1-OX40L construct, an approximately 20% to 30% reversal of the RPE-mediated suppression of PBMCs was observed. A significant difference in the abrogation of RPE-mediated suppression was not observed between the RPE transfected with the eYFP-N1-OX40L and the eYFP-C1-OX40L vectors. Interestingly, when both OX40L-C and GITR ligand (GITRL) were transfected into RPE cells, an increase in the reversal of RPE-mediated suppression was observed when compared to the reversal demonstrated with RPE transfected with OX40L-C or GITRL alone (data not shown).

#### Discussion

A significant body of evidence has supported the role of RPE cells in the maintenance of ocular immune privilege [27,31-33]. RPE cells are capable of acting as efficient APCs expressing MHC class II molecules, and exposure



**Figure 3** RT-PCR demonstrated expression of OX40L in the RNA of ARPE cells stimulated by pro-inflammatory cytokines. Lanes 1 and 2 are two different DNA ladders; lanes 3 and 4 represent two different concentrations of the same RNA from stimulated RPE cells.

to varying cytokine milieus may lead to RPE-mediated T cell activation or T cell regulation [31,34].

A number of the TNF receptor family members are thought to promote inflammatory processes via T cell co-stimulation, proliferation, and differentiation. The TNF-R/TNF- $\alpha$  interaction has been well described both in experimental animal models and patients with ocular inflammatory diseases. We previously showed that another member of the TNF receptor family, GITRL, may also play a role in ocular inflammation. Specifically, GITRL expression was able to abrogate the immunosuppressive function of RPE cells on CD3+ T cells via decreased TGF-beta expression in RPE/CD3+ T cell co-cultures and increased production of pro-inflammatory cytokines and chemokines [30]. In patients with uveitis, GITR was found to be an activation marker for CD4+ T cells and was co-expressed with CD25 on CD4+ T cells in peripheral blood. The presence of GITR was also found to be correlated with disease course in uveitis [35].

In this study, we investigated whether OX40L, another member of the TNF family involved with inflammatory disease states, may play a role in the pathogenesis of ocular inflammation. However, current human RPE cell lines do not express OX40L at the protein level. Moreover, in unactivated human RPE cells, we did not observe OX40L at the RNA level. Nonetheless, after exposure of RPE cells to a pro-inflammatory cytokine milieu (IL-1 $\alpha$ , TNF- $\alpha$ , and IFN- $\gamma$ ), we did observe OX40L RNA expression. This represents the first demonstration of OX40L RNA in ocular tissue and suggests that OX40L expression is an inducible phenomenon in the setting of localized inflammation.


To investigate the functional role of OX40L in RPE cells, we used an *in vitro* expression approach to mimic a pro-inflammatory environment. The use of a fluorescent-tagged fusion vector allowed us to confirm the presence of OX40L transfection into RPE cells and its expression using flow cytometry. In RPE-PBMC co-culture experiments, we found that RPE cells expressing OX40L (after transfection) reversed the immunosuppression afforded by RPE cells that did not express OX40L. PBMCs rather than purified T cells were used to simulate an extracellular milieu that would more closely mimic the ocular immune environment *in vivo*. OX40L was transfected in ARPE cells, which normally do not express OX40L, to mimic the ocular inflammatory scenario.

Recent reports have shown that the OX40-OX40L interaction may play a role in a number of ocular inflammatory processes including corneal transplant rejection [21], experimental allergic conjunctivitis [20], herpes stromal keratitis [36], and uveitis. Several recent studies have described the relationship of OX40 ligand in the exacerbation and prolongation of animal models of uveitis as well as its role in augmenting the pro-inflammatory Th17 cytokine response.

Moreover, several other mechanisms by which OX40L perpetuates the inflammatory process in other biologic systems have been suggested. OX40L was recently found to inhibit the generation of IL-10 producing CD4+ type 1 regulatory T cells from naive T cells. In addition, OX40L inhibited IL-10 production in differentiated IL-10 producing regulatory T cells [37]. Other proposed functions of OX40L include its role in the T cell effector response, T cell-APC co-stimulatory signaling, the T cell-dependent humoral responses [38], and the promotion of CD4+ T cell longevity [39].

## Conclusions

The numerous mechanisms by which OX40L may activate inflammation or abrogate immunosuppression, particularly in the complex ocular immune environment in which immune privilege is critical to immunoregulation,



remain incompletely understood. We have demonstrated that a pro-inflammatory cytokine milieu is sufficient to induce RNA expression of OX40L by RPE cells, and its expression appears to be highly regulated. In addition, our co-culture experiments with activated PBMCs and transfected RPE cells demonstrate that OX40L expression is associated with dampening of RPE-mediated immunosuppression, a novel finding that warrants additional investigation. This study, along with prior work on the role of GITRL in ocular inflammation, adds to the growing evidence that other members of the TNF ligand superfamily besides TNF- $\alpha$  may play a role in ocular immunity. Further studies will be needed to determine the precise role of OX40L in human uveitis and whether modulation of OX40L pathways may have therapeutic implications.

## Methods

### Reagents, cell lines, and antibodies

The retinal pigment epithelial cells (ARPE-19 cells) were originally obtained from ATCC (Manassas, VA, USA) and were cultured in minimum essential medium (MEM) with 10% fetal bovine serum (FBS; Gemini Bioproducts, West Sacramento, CA, USA), 1X nonessential amino acids, 1X antibiotics, and 2 mM glutamine (Biosource International, Bethesda, MD, USA) at 37°C in 5% CO<sub>2</sub>. Biotinylated polyclonal goat anti-human OX40L antibody (Catalog # BAF1054) was obtained from R & D systems (Minneapolis, MN, USA). IL-1 $\alpha$ , TNF- $\alpha$ , and IFN- $\gamma$  were purchased from Biosource International (Camarillo, CA, USA). The media and FBS were purchased as lipopolysaccharide-free-grade reagents.

### Plasmid

The fusion vectors eYFP-C1 and eYFP-N1 (BD Biosciences Clontech, San Jose, CA, USA) were used so that we could readily assess the presence of stably inserted OX40L DNA. The eYFP-C1 vector attaches the tag protein eYFP to the C-terminal (intracellular) region of the inserted gene, while the eYFP-N1 vector attaches eYFP protein to the N-terminal (extracellular) region of the inserted gene.

### Cloning of full-length hOX40L cDNA

Full-length OX40L was amplified from a human HTLV-1-infected T cell lymphoma (MT-2) cell line [40] using hOX40L specific primers:

hOX40L to insert into the eYFP-C1 vector

5'-CCCAAGCTTCCATGGAAAGGGTCCAACCCC-3' (forward) and

5'-CGCGGATCCTCAAAGGACACAGAATTCACC-3' (reverse).

hOX40L to insert into eYFP-N1 vector

5'-CCCAAGCTTGCCACCATGGAAAGGGTCCAACCCC-3' (forward) and

5'-CGCGGATCCGCAAGGACACAGAATTCACCAGG-3' (reverse).

RT-PCR was performed with the one-step RT-PCR kit (Promega, Madison, WI, USA) according to the manufacturer's instructions. The amplification procedure for RT-PCR was as follows: 48°C for 75 min, 94°C for 2 min and 30 s followed by cycles of the following: 94°C for 30 s, 50°C for 45 s, and 72°C for 1 min. There was a total of 40 cycles.

Total human OX40L product (552 bp) was purified using the QIAquick PCR purification kit (Qiagen Inc., Santa Clarita, CA, USA), and full-length OX40L was subcloned in frame into either the eYFP-C1 or eYFP-N1 vector. With eYFP-C1, the gene was inserted between the eYFP coding sequence and the stop codon, and with eYFP-N1, the OX40L gene was inserted between the immediate early promoter of CMV and the eYFP coding sequence.

#### Transfection of ARPE-19 cell line

To facilitate the expression of hOX40L in RPE cells, human adult retinal pigment epithelial cell line ARPE-19 was transfected with the construct eYFP-C1-OX40L or with eYFP-N1-OX40L using the Nucleofector transfection system (Amaxa Inc, Gaithersburg, MD, USA). Prior to transfection, RPE cells were grown to 80% to 85% confluence and resuspended in 2 to 3 × 10<sup>6</sup> cells/0.1 mL Nucleofector Solution V with 2 µg of eYFP-OX40L construct. The solution was transferred to a sterile electroporation cuvette and transfected using program #T20 according to manufacturers' instructions. The transfected cells were then plated in the culture medium (MEM, 10% FBS, 1X antibiotics) on six-well culture plates.

#### Expression of OX40L

Flow cytometry was utilized to analyze the surface expression of OX40L in RPE cells and to study the OX40L expression patterns in a time trial and at varying transfection dosages. For the time trial, RPE cells (2 × 10<sup>6</sup> cells) transfected with either eYFP-C1-OX40L or eYFP-N1-OX40L for different time points (30 min, 1, 4, 6, and 24 hr) were incubated in cold PBS (pH 7.4) with 0.5% BSA with or without biotinylated anti-OX40L antibody for the detection of surface OX40L. Cells were washed and fixed with 1% paraformaldehyde, counted via the FACS Calibur flow cytometer (Becton Dickinson, San Jose, CA, USA), and analyzed using FlowJo software (TreeStar, San Jose, CA, USA). Total RNA from cultured RPE cell lines was isolated with an RNA isolation kit (RNeasy, Qiagen Inc.) using the manufacturers' instructions. Prior to RNA isolation, one cell line was stimulated for 24 h with 5.0 ng/mL of IL-1α, 5.0 ng/mL of TNF-α, and 25 ng/mL of IFN-γ. Another cultured cell line was transfected with 1 µg eYFP-OX40L construct and served as the positive control for the RT-PCR. The last cultured cell line was not treated with pro-inflammatory cytokines and represented resting RPE cells. Using 1 µg of RNA, initially, RT-PCR was used to screen for the product (OX40L). The amplification procedure was as follows: 48°C for 75 min, 94°C for 2 min and 30 s followed by cycles of the following: 94°C for 30 s, 50°C for 45 s, and 72°C for 1 min. There was a total of 40 cycles. The primers used for amplification were from different exons and their sequences were as follows:

OX40L, 5'-TCACCTACATCTGCCTGCACTTCTCTGCTCTT-3' and 5'-ATCAGTTCTCCGCCATTCACA-3'.

#### Isolation of peripheral blood mononuclear cells

Human peripheral blood mononuclear cells (PBMCs) were isolated from buffy coat obtained from normal, healthy donors (NIH blood bank) using the Ficoll gradient centrifugation.

#### RPE-PBMC co-culture studies

An RPE-PBMC co-culture system was used for the *in vitro* proliferation studies. ARPE-19 cells were grown to 80% to 85% confluence and transfected with eYFP-OX40L vector. Twelve hours after transfection, the RPE cells were lethally irradiated (9,000 rads), washed with FBS-supplemented RPMI 1640 medium and plated in 96-well flat-bottomed culture plates, using enough RPE cells to obtain complete confluence. Isolated PBMCs (2 × 10<sup>5</sup> cells/well) were co-cultured with the already plated RPE cells. To a set number of cultures, anti-CD28 antibody (2 µg/mL) and anti-CD3 antibody (2 µg/mL) were added. After 3 days of culturing at 37°C in 5% CO<sub>2</sub>, the cultures were pulsed with <sup>3</sup>H thymidine (Amersham, Buckinghamshire, UK) and then cultured for another 8 to 12 h. The cells were harvested, and uptake of <sup>3</sup>H thymidine was measured by a beta counter (Perkin Elmer Life Sciences, Waltham, MA, USA). The effect of RPE cells (± transfected with OX40L) on PBMCs (± stimulation) was expressed as delta counts per minute (Δcpm). All experiments were performed in triplicate and summarized herein.

All research conformed to the ARVO statement on human and animal research and the Declaration of Helsinki. Samples from normal donors were obtained from the National Institutes of Health blood bank after informed consent was obtained.

#### Competing interests

The authors declare that they have no competing interests.

#### Authors' contributions

MC participated in the study design and coordination, carried out the experiments, and drafted the manuscript. ZL conceived of the study and participated in the study design, various experiment portions, and in the revision of the manuscript as well as the statistical analysis. BL participated in the experiments and in the revision of the manuscript. SY participated in the various experiments, in intellectual discussion, and in the drafting/revision of the manuscript. RN participated in the design of the study, in intellectual discussion, and held a supervisory role. All authors read and approved the final manuscript.

#### Author details

<sup>1</sup>Vitreoretinal Service, Department of Ophthalmology and Visual Sciences, The University of Iowa Hospitals & Clinics, Iowa City, IA 52242, USA. <sup>2</sup>National Institute of Allergy and Infectious Diseases, National Institutes of Health, Bethesda, MD 20892, USA. <sup>3</sup>Laboratory of Immunology, National Eye Institute, National Institutes of Health, 10 Center Dr. Bldg 10, 10D45, Bethesda, MD 20892, USA. <sup>4</sup>Emory Eye Center, Section of Vitreoretinal Surgery and Disease, Atlanta, GA 30322, USA.

Received: 20 September 2012 Accepted: 1 October 2012

Published: 15 January 2013

## References

1. Baum PR, Gayle RB 3rd, Ramsdell F, Srinivasan S, Sorensen RA, Watson ML, Seldin MF, Clifford KN, Grabstein K, Alderson MR (1994) Identification of OX40 ligand and preliminary characterization of its activities on OX40 receptor. *Circ Shock* 44:30–34
2. Godfrey WR, Fagnoni FF, Harara MA, Buck D, Engleman EG (1994) Identification of a human OX-40 ligand, a costimulator of CD4+ T cells with homology to tumor necrosis factor. *J Exp Med* 180:757–762
3. Gruss HJ (1996) Molecular, structural, and biological characteristics of the tumor necrosis factor ligand superfamily. *Int J Clin Lab Res* 26:143–159
4. Miura S, Ohtani K, Numata N, Niki M, Ohbo K, Ina Y, Gojobori T, Tanaka Y, Tozawa H, Nakamura M (1991) Molecular cloning and characterization of a novel glycoprotein, gp34, that is specifically induced by the human T-cell leukemia virus type I transactivator p40tax. *Mol Cell Biol* 11:1313–1325
5. Murata K, Ishii N, Takano H, Miura S, Ndhlovu LC, Nose M, Noda T, Sugamura K (2000) Impairment of antigen-presenting cell function in mice lacking expression of OX40 ligand. *J Exp Med* 191:365–374
6. Chen AI, McAdam AJ, Buhlmann JE, Scott S, Lupher ML Jr, Greenfield EA, Baum PR, Fanslow WC, Calderhead DM, Freeman GJ, Sharpe AH (1999) OX40-ligand has a critical costimulatory role in dendritic cell: T cell interactions. *Immunity* 11:689–698
7. Gruss HJ, Dower SK (1995) Tumor necrosis factor ligand superfamily: involvement in the pathology of malignant lymphomas. *Blood* 85:3378–3404
8. Murata K, Nose M, Ndhlovu LC, Sato T, Sugamura K, Ishii N (2002) Constitutive OX40/OX40 ligand interaction induces autoimmune-like diseases. *J Immunol* 169:4628–4636
9. Watts TH, DeBenedette MA (1999) T cell co-stimulatory molecules other than CD28. *Curr Opin Immunol* 11:286–293
10. Vetto JT, Lum S, Morris A, Sicotte M, Davis J, Lemon M, Weinberg A (1997) Presence of the T-cell activation marker OX-40 on tumor infiltrating lymphocytes and draining lymph node cells from patients with melanoma and head and neck cancers. *Am J Surg* 174:258–265
11. Weinberg AD, Bourdette DN, Sullivan TJ, Lemon M, Wallin JJ, Maziarz R, Davey M, Palida F, Godfrey W, Engleman E, Fulton RJ, Offner H, Vandenberg AA (1996) Selective depletion of myelin-reactive T cells with the anti-OX-40 antibody ameliorates autoimmune encephalomyelitis. *Nat Med* 2:183–189
12. Stuber E, Neurath M, Calderhead D, Fell HP, Strober W (1995) Cross-linking of OX40 ligand, a member of the TNF/NGF cytokine family, induces proliferation and differentiation in murine splenic B cells. *Immunity* 2:507–521
13. Kato H, Kojima H, Ishii N, Hase H, Imai Y, Fujibayashi T, Sugamura K, Kobata T (2004) Essential role of OX40L on B cells in persistent alloantibody production following repeated alloimmunizations. *J Clin Immunol* 24:237–248
14. Ndhlovu LC, Ishii N, Murata K, Sato T, Sugamura K (2001) Critical involvement of OX40 ligand signals in the T cell priming events during experimental autoimmune encephalomyelitis. *J Immunol* 167:2991–2999
15. Hoshino A, Tanaka Y, Akiba H, Asakura Y, Mita Y, Sakurai T, Takaoka A, Nakaïke S, Ishii N, Sugamura K, Yagita H, Okumura K (2003) Critical role for OX40 ligand in the development of pathogenic Th2 cells in a murine model of asthma. *Eur J Immunol* 33:861–869
16. Ohshima Y, Tanaka Y, Tozawa H, Takahashi Y, Maliszewski C, Delespesse G (1997) Expression and function of OX40 ligand on human dendritic cells. *J Immunol* 159:3838–3848
17. Brocker T, Gulbranson-Judge A, Flynn S, Riedinger M, Raykundalia C, Lane P (1999) CD4 T cell traffic control: *in vivo* evidence that ligation of OX40 on CD4 T cells by OX40-ligand expressed on dendritic cells leads to the accumulation of CD4 T cells in B follicles. *Eur J Immunol* 29:1610–1616
18. Weinberg AD, Wegmann KW, Funatake C, Whitham RH (1999) Blocking OX-40/OX-40 ligand interaction *in vitro* and *in vivo* leads to decreased T cell function and amelioration of experimental allergic encephalomyelitis. *J Immunol* 162:1818–1826
19. Imura A, Hori T, Imada K, Ishikawa T, Tanaka Y, Maeda M, Imamura S, Uchiyama T (1996) The human OX40/gp34 system directly mediates adhesion of activated T cells to vascular endothelial cells. *J Exp Med* 183:2185–2195
20. Fukushima A, Yamaguchi T, Ishida W, Fukata K, Yagita H, Ueno H (2006) Roles of OX40 in the development of murine experimental allergic conjunctivitis: exacerbation and attenuation by stimulation and blocking of OX40. *Invest Ophthalmol Vis Sci* 47:657–663
21. Hattori T, Usui Y, Okunuki Y, Sonoda Y, Usui M, Takada E, Mizuguchi J, Yagita H, Okumura K, Akiba H, Takeuchi M (2007) Blockade of the OX40 ligand prolongs corneal allograft survival. *Eur J Immunol* 37:3597–3604
22. Zhang Z, Zhong W, Hinrichs D, Wu X, Weinberg A, Hall M, Spencer D, Wegmann K, Rosenbaum JT (2010) Activation of OX40 augments Th17 cytokine expression and antigen-specific uveitis. *Am J Pathol* 177:2912–2920
23. Wu X, Rosenbaum JT, Adamus G, Zhang GL, Duan J, Weinberg A, Zhang Z (2011) Activation of OX40 prolongs and exacerbates autoimmune experimental uveitis. *Invest Ophthalmol Vis Sci* 52:8520–8526
24. Niederkorn JY (2006) See no evil, hear no evil, do no evil: the lessons of immune privilege. *Nat Immunol* 7:354–359
25. Niederkorn JY (2005) A renewed appreciation for ocular immune privilege. *Arch Soc Exp Ophthalmol* 80:437–438, 9–40
26. Niederkorn JY (1990) Immune privilege and immune regulation in the eye. *Adv Immunol* 48:191–226
27. Kaestel CG, Lovato P, Odum N, Nissen MH, Ropke C (2005) The immune privilege of the eye: human retinal pigment epithelial cells selectively modulate T-cell activation *in vitro*. *Curr Eye Res* 30:375–383
28. Jorgensen A, Wiencke AK, la Cour M, Kaestel CG, Madsen HO, Hamann S, Lui GM, Scherfig E, Prause JU, Svejgaard A, Odum N, Nissen MH, Röpke C (1998) Human retinal pigment epithelial cell-induced apoptosis in activated T cells. *Invest Ophthalmol Vis Sci* 39:1590–1599
29. Kaestel CG, Jorgensen A, Nielsen M, Eriksen KW, Odum N, Holst Nissen M, Röpke C (2002) Human retinal pigment epithelial cells inhibit proliferation and IL2R expression of activated T cells. *Exp Eye Res* 74:627–637
30. Mahesh SP, Li Z, Liu B, Fariss RN, Nussenblatt RB (2006) Expression of GITR ligand abrogates immunosuppressive function of ocular tissue and differentially modulates inflammatory cytokines and chemokines. *Eur J Immunol* 36:2128–2138
31. Sun D, Enzmann V, Lei S, Sun SL, Kaplan HJ, Shao H (2003) Retinal pigment epithelial cells activate uveitogenic T cells when they express high levels of MHC class II molecules, but inhibit T cell activation when they express restricted levels. *J Neuroimmunol* 144:1–8
32. Ishida K, Panjwani N, Cao Z, Streilein JW (2003) Participation of pigment epithelium in ocular immune privilege. 3. Epithelia cultured from iris, ciliary body, and retina suppress T-cell activation by partially non-overlapping mechanisms. *Ocul Immunol Inflamm* 11:91–105
33. Gregerson DS, Heuss ND, Lew KL, McPherson SW, Ferrington DA (2007) Interaction of retinal pigmented epithelial cells and CD4 T cells leads to T-cell anergy. *Invest Ophthalmol Vis Sci* 48:4654–4663
34. Percopo CM, Hooks JJ, Shinohara T, Caspi R, Detrick B (1990) Cytokine-mediated activation of a neuronal retinal resident cell provokes antigen presentation. *J Immunol* 145:4101–4107
35. Li Z, Mahesh SP, Kim BJ, Buggage RR, Nussenblatt RB (2003) Expression of glucocorticoid induced TNF receptor family related protein (GITR) on peripheral T cells from normal human donors and patients with non-infectious uveitis. *J Autoimmun* 21:83–92
36. Lepisto AJ, Xu M, Yagita H, Weinberg AD, Hendricks RL (2007) Expression and function of the OX40/OX40L costimulatory pair during herpes stromal keratitis. *J Leukoc Biol* 81:766–774
37. Ito T, Wang YH, Duramad O, Hanabuchi S, Perng OA, Gilliet M, Qin FX, Liu YJ (2006) OX40 ligand shuts down IL-10-producing regulatory T cells. *Proc Natl Acad Sci USA* 103:13138–13143
38. Stuber E, Strober W (1996) The T cell-B cell interaction via OX40-OX40L is necessary for the T cell-dependent humoral immune response. *J Exp Med* 183:979–989
39. Soroosh P, Ine S, Sugamura K, Ishii N (2006) OX40-OX40 ligand interaction through T cell-T cell contact contributes to CD4 T cell longevity. *J Immunol* 176:5975–5987
40. Harada S, Koyanagi Y, Yamamoto N (1985) Infection of HTLV-III/LAV in HTLV-I-carrying cells MT-2 and MT-4 and application in a plaque assay. *Science* 229:563–566

doi:10.1186/1869-5760-3-12

Cite this article as: Cunningham et al.: OX40 ligand expression abrogates the immunosuppressive function of retinal pigment epithelium. *Journal of Ophthalmic Inflammation and Infection* 2013 3:12.